

conditioned media from *Bifidobacterium infantis* prevented a reduction in transepithelial resistance of intestinal epithelial cells and reduced ileal and colonic permeability in IL-10-deficient mice.¹⁰ Another study demonstrated that prior exposure of an intestinal epithelial cell line to viable *L. acidophilus*, but not heat-inactivated *L. acidophilus*, limited the adverse effects induced by an *Escherichia coli* strain, such as a decline in transepithelial resistance, increased epithelial permeability and physiological dysfunction. Additionally, viable *L. acidophilus* reduced phosphorylation of tight junction proteins induced by *E. coli*.⁶ Two proteins, p40 and p75, were purified from the culture supernatant of *L. rhamnosus* LGG, and both proteins prevented TNF-induced apoptosis and intestinal barrier disruption in colonic epithelial cells.¹¹

The effects of probiotics and their effector molecules on barrier function in clinical trials are sparse. In one study, the efficacy of a placebo, VSL#3 and probiotic-derived sonicates, was studied for modulation of intestinal permeability in patients in the intensive care unit with multiple organ dysfunction syndrome.¹² Patients that received VSL#3 exhibited a larger increase in serum IgA and IgG concentrations than the patients who received placebo or sonicates. However, no significant difference in intestinal permeability was observed between the patients who received VSL#3, probiotic sonicates and placebo.¹² In another study, a short-term improvement in mucosal barrier function in patients with IBS was observed after administration of a probiotic fermented milk (*Streptococcus thermophilus*, *Lactobacillus bulgaricus*, *L. acidophilus* and *Bifidobacterium longum*) compared to a milk beverage containing no bacteria.¹³ These studies emphasize the importance of strain selection and viability. In addition, as noted in other studies with prebiotics and probiotics, effects on gut barrier function varied between strains and effector molecules.^{6,10,14} It is clear that probiotic immunomodulatory components consist of a varied array of effector molecules (Table 2), including surface layer proteins, cell wall polysaccharides, adhesins, teichoic acids and heat shock proteins. Other immunomodulatory components have been understudied or remain to be characterized. While in vitro work to identify important candidate effector molecules (Table 2) continues, the effects on mucosal immunity in human intervention studies still needs to be determined.

Live bacteria, dead bacteria or bacterial supernatants. Effects on host gene expression and mucosal immune responses. Studies have been performed using live, dead and bacterial supernatants to assess the mucosal immune response. In particular, probiotic *L. plantarum* WCFS1 has been studied using in situ-based -omic models in the GIT. Gene expression of beneficial bacteria¹⁵ and the mammalian host^{16,17} has been studied in parallel. Gene expression profiles in ceca of germ-free mice fed a standard low fat or western diet were investigated to determine the effects of host diet on bacterial (probiotic) gene expression in the GIT.¹⁵ The western diet was high in simple sugars and fat, and the standard diet (low fat diet) was high in complex plant polysaccharides.¹⁵ In addition to transcriptomics studies, colonization levels of *L. plantarum* were measured in mice fed both diet types, and was determined to be ten times higher in mice receiving the standard feed compared to the western diet. Gene expression was

Table 1. Key Questions require further study and discussion

- What are the specific probiotic components that mediate immunomodulation?
- What other microbes, perhaps as yet undiscovered or under studied exhibit immunomodulation properties?
- What is considered healthy and how do we measure health benefits in human trials and animal studies?
- What are the factors influencing responses in clinical trials?
- Which biomarkers and immune parameters should we use in probiotic and prebiotic intervention studies in humans?
- What are the systemic effects of prebiotics on the gut microbiota and direct effects on the gut mucosa?
- What are the best target groups to use for human trials?
- How comparable are in vitro, versus in vivo and ex vivo studies?
- How comparable are results from animal models to human clinical trials?

compared to a reference group of differentially expressed genes, namely mid-logarithmic *L. plantarum* grown in vitro in MRS, chemically defined medium and chow medium. The main results from this study indicated that bacterial genes involved in carbohydrate metabolism and cell surface functions were upregulated in mice fed the standard diet compared to the culture media. In addition, it was determined that the western diet resulted in a nutritionally restricted, growth-limiting environment for *L. plantarum* in the distal gut.¹⁵ In mice fed both diets, gene sets encoding cell surface-related functions were differentially expressed, which included genes involved in D-alanylation of LTA, a probiotic effector molecule (Table 2). The authors suggested that the probiotic bacteria modified its gene expression to reduce levels of LTA on the cell surface, as a means to restrict exposure to components of the host immune system in the GIT of mice fed both diet types.

The growth phase and viability of the probiotic cultures is also a factor that varies between reported experiments. For example, in a study by van Baarlen and co-workers,¹⁷ differences in gene expression in the duodenum of healthy humans was observed depending on the growth phase and viability of the probiotic *L. plantarum* WCFS1. Individuals were administered logarithmic, stationary or dead cells and samples were taken from the duodenum for analysis after 6 hours. Results from this small randomized, double-blind, placebo-controlled crossover study identified gene expression patterns at the mucosal surface and within cellular pathways that were related to immune tolerance. While all three conditions resulted in the induction of genes involved in the immune response, differences in the type of induction and number of genes induced were observed between logarithmic and stationary phase cells. Logarithmic phase cells induced a response targeted towards metabolic functions such as nucleic acid metabolism and cytoplasm organization, however stationary phase cells resulted in induction of genes such as the NFκB and JUN transcription factors involved in the establishment of immune tolerance.¹⁷ Interestingly, the dead cells (heat killed stationary phase) induced the highest fold changes, and the mucosal responses to dead and stationary bacteria were comparable to

Table 2. Potential effector molecules derived from probiotics

Bacteria	Molecule	Effect	References
<i>Lactobacillus reuteri</i>	Secreted factors by <i>L. reuteri</i> biofilms	Suppression of human TNF production by LPS-activated monocyotoid cells	54
<i>Lactobacillus reuteri</i>	Secretion of reuterin (antimicrobial glycerol derivative) by <i>L. reuteri</i> biofilms	Secretion of reuterin by biofilms demonstrated	54
Intestinal Bacteria	Butyrate (a short chain fatty acid)	Increased intestinal barrier function in Caco2 cell lines	55
<i>Lactobacillus acidophilus</i> NCFM	S layer protein	DC-SIGN ligand involved in the modulation of DCs and T cells functions	56
<i>Faecalibacterium prausnitzii</i>	<i>F. prausnitzii</i> supernatant	Reduced the severity of TNBS colitis and tended to correct the dysbiosis associated with TNBS colitis	57
<i>Lactobacillus casei</i> Shirota	Cell wall polysaccharide moiety	Inhibit macrophage and splenocyte activity	58
<i>Bifidobacteria infantis</i>	<i>B. infantis</i> conditioned medium	Reduced colonic permeability in mice and attenuated inflammation in IL-10-deficient mice	10
<i>Lactobacillus helveticus</i>	S layer protein	Inhibition of <i>Escherichia coli</i> 0157:H7 adherence to intestinal cell lines	59
<i>Lactobacillus rhamnosus</i> GG	Modified teichoic acids	Increased sensitivity to gastric juice and human beta-defensin-2 but no difference in immunomodulation	60
<i>L. rhamnosus</i> GG	P40 and P75 secreted proteins	Prevention of cytokine-induced apoptosis in human and mouse intestinal epithelial cells	11
<i>Lactobacillus salivarius</i> UCC118	Secretion of bacteriocin Abp118	Protected mice against infection with the pathogen <i>Listeria monocytogenes</i>	61
<i>Escherichia coli</i> Nissle 1917	Flagellin	Induction of beta-defensin-2 in intestinal cell lines	62
<i>Lactobacillus johnsonii</i> La1	GroEL	Bind to components of the gastrointestinal mucosa and stimulates interleukin-8 secretion in macrophages and HT29 cells in a CD14-dependent mechanism	63
<i>Lactobacillus delbrueckii</i> ssp. <i>bulgaricus</i> OLL1073R	Extracellular polysaccharides (EPS)	Stimulation of mouse splenocytes and significant increase in interferon-gamma production	64
<i>Lactobacillus plantarum</i> NCIMB8826	Modified teichoic acids	Protective against colitis	23
<i>L. johnsonii</i> NCC533	Adhesin; elongation factor Tu (EF-Tu)	Mucin binding, proinflammatory response in the presence of soluble CD14	65

each other but different to the response from logarithmic bacteria. These studies demonstrated the importance of reference sets of gene expression data used for comparisons in vivo, in addition to diet, growth media and growth phase of probiotics. Factors such as the stability of each probiotic strain, expression and stability of effector molecule, growth phase of the probiotic strain, site of action of probiotic-derived effector molecules should be considered for in vivo studies. Finally, can immunomodulation in the laboratory be connected with health benefits in humans and animals?

Prebiotics and mucosal immunity. Prebiotic benefits include modulation of the gut microbiota, increased mineral adsorption, modulation of lipid metabolism and inhibition of pathogens.¹⁸ While studies of prebiotics have documented their effects on mucosal immunity, the majority of these reports included animal models, with relatively few human studies published to date. Some published studies have included other factors such as antioxidants and vitamins, which may also contribute to mucosal immunity. In studies with synbiotics, it is difficult to distinguish the effects of prebiotics from that of probiotics.

A few studies examined immunomodulation by dietary prebiotics (reviewed in ref. 19). Prebiotic supplementation increased fecal secretory IgA and postnatal immune development in infants.²⁰ In disease states, moderate effects were seen in patients with Crohn's disease,²¹ but results were encouraging enough to warrant further exploration. Because prebiotic compounds per definition modulate the composition of the gut microbiota, it is difficult to know whether these effects are direct or are the consequences of a shift of certain bacterial groups. Clinical trials have included an analysis of microbial composition and associated effects on short chain fatty acids (SCFA). Products of bacteria that have been reported to show anti-inflammatory properties include SCFAs (Table 2),²² and other bacterial metabolites that could be responsible for the modulation of the immune system, remain to be characterized.

Additional animal and human studies are needed, as there is no clear consensus on the effects of prebiotics on mucosal immunity. Points to consider include the nature of prebiotics and consequent indirect interactions such as system-wide changes of gut microbiota and the subsequent effects on mucosal immunity

and/or health benefits and direct interactions of prebiotics with the mucosa. The challenge is determining these effects on health and designing better trials including the use of biomarkers (discussed below).

Can we Extrapolate Immunomodulatory Effects of Probiotics and Prebiotics in vitro to Humans?

Factors influencing the immune response in vivo and in human clinical trials. Extrapolation of immunomodulatory effects found in the laboratory and in animal studies with outcomes in human trials presents a difficult challenge. Immunomodulatory effects conferred by *L. plantarum* WCFS1 in vitro,²³ in animal models,^{16,23} but also in humans^{24,17} highlights the difficulties of comparing similar effects by a single strain in different contexts. A few examples show correlations between in vitro and in vivo immunomodulatory properties of lactic acid bacteria.^{23,25,26} The study by Foligne et al.²⁵ for instance, demonstrated the possibility of translating results from an in vitro assay with peripheral blood mononuclear cells (PBMCs) obtained from different donors, to a murine model of colitis using the IL-10/IL-12 ratio. Generally, the discrepancies between in vitro and in vivo results observed in published trials can be partly explained by the host contribution (genetic factors, different baseline immune functions between individuals, microbiome diversity, differences in the body sites targeted, intra-person variation) as well as environmental factors (diet, stress etc.,) partially controlled by each individual. Different factors can affect outcomes in studies reporting immunomodulatory effects of probiotic or prebiotic interventions in vivo.

Variation between individuals and effects on host response.

Genetic factors. Genetic variability strongly influences the immune response of the host. Specific polymorphisms in genes encoding antigens (human leukocytes²⁷) or proteins involved in the immune response can influence quantities of cytokines²⁸ in a healthy population or disease state. A growing body of evidence indicates that individuals respond to bacterial signals (TLR ligands) differently due to single nucleotide polymorphisms (SNPs) within TLR genes.²⁹ A frameshift mutation in NOD2 is a strong risk factor in patients with inflammatory bowel disease (Crohn's disease), and this evidence supports the importance of SNPs and inter-individual differences in TLR signaling.^{30,31}

Microbiome. Preliminary observations that all vertebrates have a microbiome that co-evolved with the host, resulted in the hypothesis that the adaptive immune system (memory-based) system evolved in the vertebrate lineage because of the intimate co-existence with complex communities of beneficial microbes.³² Whether this is true or not still needs to be demonstrated, but it certainly highlights the complex links between our microbiome and our immune system. Because we are in the early stages of characterizing the phylogenetic or functional core microbiomes in human and animal models, it is difficult to fully understand relationships between the host microbiota and immune system. However, the lack of microbes such as *Faecalibacterium prausnitzii* in inflammatory bowel disease (Crohn's disease) has been demonstrated by different studies.³³ Baseline immune functions

and microbiomes may differ and depend on each other so that aggregate responses and disease susceptibilities in individuals may represent a combination of microbial composition, microbial functions and host genotypes.

Influence of environmental factors controlled by the host such as diet. The gut microbiota is a dynamic environment, and changes induced by the diet, the uptake of antibiotics or even physiologic stresses may induce rapid changes in microbial composition and functions,³⁴ with a concomitant impact on the host immune response. Rapid changes in microbial composition have been well documented with a high fat diet.³⁵ These results should encourage investigators to control diets of subjects and support the use of standardized diets when measuring the host immune response. Dietary components in the diet may have a direct effect on the response of the immune system and vitamins, for example, may have an effect on cytokine production.³⁶ To make the story more complex, effects of dietary components may be partly determined by genetic polymorphisms of cytokines such as TNF.³⁷

Consideration of body site contribution. Small intestine versus colon. Immune parameters measured in different body sites will certainly be different and may have an impact of results and outcomes of studies with probiotics and prebiotics. In the gut, the colon has been more extensively investigated than the small intestine or other mucosal sites of the gastrointestinal tract such mouth or stomach,³⁸ and certainly merits further investigation. As the composition of the mucus layer and mucus barrier is different at these sites, the contribution of the mucus layer and its interaction with probiotics and prebiotics or other members of the gut microbiota should be considered and investigated further.

Which biomarkers and immune parameters shall we use in probiotic/prebiotic intervention in humans? No universal biomarkers have been identified to assess how probiotics and prebiotics impact mucosal immunity. However, specific biomarkers can be defined for certain populations and disease states, and are summarized in Table 3. Measuring these parameters is still challenging in healthy people, and limited by the ability to analyze markers only in blood, saliva, fecal samples or urine. In an experimental setting with disease patients, it is usually possible to get biopsies from the gut. New techniques and potential biomarkers are still emerging and may offer promising alternatives in the future. For instance, recent discoveries on the role of C-reactive protein (CRP) isoforms and their role in the inflammation could facilitate an improved understanding of the importance of different isoforms.³⁹ Metabolites that distinguish patients with IBD, IBS or other gut diseases have emerged.^{39,40} Choices of immune markers will certainly depend on the population studies and diseases studied in probiotic and prebiotic interventions.

Consideration of interventional trials with probiotics and prebiotics. Choices of animal models. Different animal models have been used to study the effects of probiotics and prebiotics on mucosal immunity. Effects in mouse models of IBD have included IL-10 knockout mice, trinitrobenzene sulfonic acid (TNBS)- or dextran sodium sulphate (DSS)-induced colitis. Based on the characterization of the microbiota in different knockout mice and wild-type mice using genotyping and 16S rRNA gene analyses, large differences in the composition of the microbiota can

Table 3. Possible immune biomarkers and parameters for assessment of mucosal and systemic immunity in vivo in response to probiotic or prebiotic interventions

Marker	Advantages	Drawbacks	References
Cytokines (TNF, IL6, IL10, IL12) FoxP3	Easy to measure	Level in blood may not reflect levels in other body sites	66
CRP and acute phase reactants	Easy to measure Different conformational change	Measure other outcome	39
Antibody (IgA)	Easy to measure	Not helpful without immune challenge	67
Gene expression profile in tissues and peripheral blood (targeted vs. global)	Extensive survey of gene activated	Difficulties for obtaining tissues in healthy humans	17, 24
Proteomics and metabolomics in urine and blood and fecal water	Extensive survey of metabolites produced	Difficult to analyze Lack of references	44, 68, 69
Calprotectin	Easy to measure in stool	Standardization necessary	70, 71, 67

be observed in different animal models,⁴¹ as well as between animal models obtained from different vendors.⁴² Differences in microbial composition between C57BL/6 mice obtained from the Jackson laboratory and Taconic laboratory was investigated recently using a high density microarray (PhyloChip). These mice differed with respect to the relative proportions of Th17 cells, cell that contribute to inflammation and may be mediators in immune protection and immunopathology. Comparative analyses of 766 bacterial taxa detected in both groups of mice demonstrated large differences in the relative abundance of more than half of the taxa, with two taxa (*Lactobacillus murinus* and segmented filamentous bacterium [SFB]) being more than 25-fold more abundant in Taconic mice. The Taconic mice in this study had elevated proportions of Th17 cells.⁴³ Most importantly, it was demonstrated that colonization of mice housed in the Jackson laboratory with one SFB species (*Arthromitus* spp.), was sufficient to induce the appearance of Th17 cells derived from a CD4⁺ T helper cell lineage.⁴³ Furthermore, this colonization was correlated with enhanced expression of genes associated with inflammation and antimicrobial defenses, resulting in enhanced resistance to the murine intestinal pathogen *Citrobacter rodentium*. Thus, characterizing the microbiomes in animals at sites where immune parameters will be measured, before intervention, may provide further insights into the possible roles of certain microbes and outcomes with probiotics or prebiotics. Murine models with “humanized” microbiomes offer promising systems for evaluation of microbial effects on host immune responses. Thus, humanized microbiome murine models (for example with a baby microbiota)⁴⁴ have similarities with that of formula-fed neonates.⁴⁵ However, the relevance of these humanized rodent models to human diseases remains to be seen.

- **Response to vaccine challenges.** Vaccine challenges are particularly useful to assess the efficacy of probiotics that may enhance the immune status in a healthy population. Probiotics enhance the immunogenicity of several vaccines including rotavirus, influenza, poliovirus, hepatitis B and pneumococcus.⁴⁶⁻⁴⁸ Measurement of antibody responses to vaccines is straightforward, and such human studies may provide excellent opportunities to evaluate the immunologic efficacy of probiotics.

- **Non-responders.** Separating responders from non-responders may facilitate interpretations of effects of the probiotics or

prebiotics on the immune system. Only a few in vivo trials have been performed.⁴⁹

- **Influence of age, gender, diet.** Because immune parameters differ according to age, gender, health status, activity and dietary habits, human studies should use crossover placebo-controlled design while trying to match and compare different groups.

- **Core or functional microbiomes.** In the future, integrated approaches using microbiology, genomics, transcriptomics and proteomics tools would certainly highlight the mechanisms and the microbes that may influence the immune system.⁵⁰ It is still debatable whether a core microbiome exists at the phylogenetic level⁵¹ or functional level.⁵² However, recent studies, such as those by Gordon and colleagues, suggest the existence of a functional core at the level of shared gene families, which suggests that different species could fulfill similar functional roles in the GIT. As pointed out by Shop et al.⁵³ however, an interesting follow-up question regarding the outcome of this study is whether the lack of shared phylotypes is still as pronounced if rare populations are taken into account.

Summary and Conclusions

Research scientists face numerous challenges including the demonstration and articulation of specific beneficial effects and associated mechanisms by probiotics and prebiotics. A clear definition of human health, if one exists, is a central consideration. What is considered “healthy” as a baseline condition may vary depending on age, gender, ethnicity, diet and numerous environmental factors? Although we have some answers with certain parameters (leukocytes, cytokines), baseline parameters including quantitative “normal” ranges of immune biomarkers remain to be defined. Non-invasive techniques that evaluate blood, fecal water or urine metabolites need to be further developed. Many studies are needed to understand various metabolites and how they may provide measurable standards for interventional studies. Additionally, the involvement of neuro-immune interactions in the GIT is an understudied area of research with respect to probiotics and prebiotics.

The difficulty to establish health claims such as “strengthen the immune response” for probiotics has been illustrated recently

in Europe by the absence of health claims related to the stimulation of the immune system authorized for probiotic foods by the European Food Safety Authority (http://www.isapp.net/docs/ISAPP_responds_to_EFSA_oct09.pdf). A growing body of evidence has documented the beneficial effects of probiotics and prebiotics in disease treatment and management studies. However, long-term studies are still needed to identify their prophylactic effects towards inflammatory disorders in different populations.

Before using probiotics in interventional studies, many factors remain to be considered. For example, viable cells are generally more effective at stimulating adaptive immunity, and the method of cell killing should be considered if nonviable cells are used. If cell supernatants are used, the active component(s) should be purified and the stability and physiologic effects of these compounds must be considered. Doses of bacteria and growth phase at time of harvest are additional considerations in tandem with

traditional methods of determining strain robustness or functional effects. On the host side, genetic influences, site of action, the mucus barrier, route of administration, diet and microbial composition of the host contribute to results of interventional trials published to date. In future trials, individual factors that potentially affect the efficacy of probiotics and prebiotics must be addressed. Continuing advancement in technologies, knowledge of the immune system, gut microbiota and improved biomarkers are essential to making human interventional studies with probiotics and prebiotics successful.

Acknowledgements

The authors would like to thank all participants of the discussion group and in particular the investigators who shared their presentations from the meeting: Peter Bron, Michael Hsieh, Flavia Indrio, Bob Rastall, Karen Madsen and Fang Yan. S.O.F. would like to thank Todd R. Klaenhammer.

References

1. Lebeer S, Vanderleyden J, De Keersmaecker SC. Host interactions of probiotic bacterial surface molecules: comparison with commensals and pathogens. *Nat Rev Microbiol* 8:171-84.
2. Netea MG, van de Veerdonk FL, Kullberg BJ, Van der Meer JW, Joosten LA. The role of NLRs and TLRs in the activation of the inflammasome. *Expert Opin Biol Ther* 2008; 8:1867-72.
3. FAO/WHO. Guidelines for the Evaluation of Probiotics in Food. London Ontario, Canada 2002.
4. Garcia-Lafuente A, Antolin M, Guarner F, Crespo E, Malagelada JR. Modulation of colonic barrier function by the composition of the commensal flora in the rat. *Gut* 2001; 48:503-7.
5. Madsen K, Cornish A, Soper P, McKaigney C, Jijon H, Yachimec C, et al. Probiotic bacteria enhance murine and human intestinal epithelial barrier function. *Gastroenterology* 2001; 121:580-91.
6. Resta-Lenert S, Barrett KE. Live probiotics protect intestinal epithelial cells from the effects of infection with enteroinvasive *Escherichia coli* (EIEC). *Gut* 2003; 52:988-97.
7. Zyrek AA, Cichon C, Helms S, Enders C, Sonnenborn U, Schmidt MA. Molecular mechanisms underlying the probiotic effects of *Escherichia coli* Nissle 1917 involve ZO-2 and PKCzeta redistribution resulting in tight junction and epithelial barrier repair. *Cell Microbiol* 2007; 9:804-16.
8. Mennigen R, Nolte K, Rijcken E, Utech M, Loeffler B, Senniger N, et al. Probiotic mixture VSL#3 protects the epithelial barrier by maintaining tight junction protein expression and preventing apoptosis in a murine model of colitis. *Am J Physiol Gastrointest Liver Physiol* 2009; 296:1140-9.
9. Rakoff-Nahoum S, Paglino J, Eslami-Varzaneh F, Edberg S, Medzhitov R. Recognition of commensal microflora by toll-like receptors is required for intestinal homeostasis. *Cell* 2004; 118:229-41.
10. Ewaschuk JB, Diaz H, Meddings L, Diederichs B, Dmytrash A, Backer J, et al. Secreted bioactive factors from *Bifidobacterium infantis* enhance epithelial cell barrier function. *Am J Physiol Gastrointest Liver Physiol* 2008; 295:1025-34.
11. Yan F, Cao H, Cover TL, Whitehead R, Washington MK, Polk DB. Soluble proteins produced by probiotic bacteria regulate intestinal epithelial cell survival and growth. *Gastroenterology* 2007; 132:562-75.
12. Alberda C, Gramlich L, Meddings J, Field C, McCargar L, Kutsogiannis D, et al. Effects of probiotic therapy in critically ill patients: a randomized, double-blind, placebo-controlled trial. *Am J Clin Nutr* 2007; 85:816-23.
13. Zeng J, Li YQ, Zuo XL, Zhen YB, Yang J, Liu CH. Clinical trial: effect of active lactic acid bacteria on mucosal barrier function in patients with diarrhoea-predominant irritable bowel syndrome. *Aliment Pharmacol Ther* 2008; 28:994-1002.
14. Nissen L, Chingwaru W, Sgorbati B, Biavati B, Cencic A. Gut health promoting activity of new putative probiotic/protective *Lactobacillus* spp. strains: a functional study in the small intestinal cell model. *Int J Food Microbiol* 2009; 135:288-94.
15. Marco ML, Peters TH, Bongers RS, Molenaar D, van Hemert S, Sonnenburg JL, et al. Lifestyle of *Lactobacillus plantarum* in the mouse caecum. *Environ Microbiol* 2009; 11:2747-57.
16. Marco ML, Bongers RS, de Vos WM, Kleerebezem M. Spatial and temporal expression of *Lactobacillus plantarum* genes in the gastrointestinal tracts of mice. *Appl Environ Microbiol* 2007; 73:124-32.
17. van Baarlen P, Troost FJ, van Hemert S, van der Meer C, de Vos WM, de Groot PJ, et al. Differential NFkappaB pathways induction by *Lactobacillus plantarum* in the duodenum of healthy humans correlating with immune tolerance. *Proc Natl Acad Sci USA* 2009; 106:2371-6.
18. Steed H, Macfarlane S. Mechanisms of Prebiotic Impact on Health. In: DC, Rastall RA, eds. *Prebiotics and Probiotics Science and Technology*. New York: Springer 2009.
19. Seifert S, Watzl B. Inulin and oligofructose: review of experimental data on immune modulation. *J Nutr* 2007; 137:2563-7.
20. Bakker-Zierikzee AM, Tol EA, Kroes H, Alles MS, Kok FJ, Bindels JG. Faecal SIgA secretion in infants fed on pre- or probiotic infant formula. *Pediatr Allergy Immunol* 2006; 17:134-40.
21. Lindsay JO, Whelan K, Stagg AJ, Gobin P, Al-Hassi HO, Rayment N, et al. Clinical, microbiological, and immunological effects of fructo-oligosaccharide in patients with Crohn's disease. *Gut* 2006; 55:348-55.
22. Maslowski KM, Vieira AT, Ng A, Kranich J, Sierro F, Yu D, et al. Regulation of inflammatory responses by gut microbiota and chemoattractant receptor GPR43. *Nature* 2009; 461:1282-6.
23. Granelle C, Nutten S, Palumbo E, Morath S, Hermann C, Dewulf J, et al. Enhanced antiinflammatory capacity of a *Lactobacillus plantarum* mutant synthesizing modified teichoic acids. *Proc Natl Acad Sci USA* 2005; 102:10321-6.
24. Troost FJ, van Baarlen P, Lindsey P, Kodde A, de Vos WM, Kleerebezem M, et al. Identification of the transcriptional response of human intestinal mucosa to *Lactobacillus plantarum* WCFS1 in vivo. *BMC Genomics* 2008; 9:374.
25. Foligne B, Nutten S, Granelle C, Dennin V, Goudercourt D, Poiret S, et al. Correlation between in vitro and in vivo immunomodulatory properties of lactic acid bacteria. *World J Gastroenterol* 2007; 13:236-43.
26. Pena JA, Rogers AB, Ge Z, Ng V, Li SY, Fox JG, et al. Probiotic *Lactobacillus* spp. diminish *Helicobacter hepaticus*-induced inflammatory bowel disease in interleukin-10-deficient mice. *Infect Immun* 2005; 73:912-20.
27. Howell WM, Calder PC, Grimble RF. Gene polymorphisms, inflammatory diseases and cancer. *Proc Nutr Soc* 2002; 61:447-56.
28. Yaqoob P, Newsholme EA, Calder PC. Comparison of cytokine production in cultures of whole human blood and purified mononuclear cells. *Cytokine* 1999; 11:600-5.
29. Schroder NW, Schumann RR. Single nucleotide polymorphisms of Toll-like receptors and susceptibility to infectious disease. *Lancet Infect Dis* 2005; 5:156-64.
30. Ogura Y, Bonen DK, Inohara N, Nicolae DL, Chen FF, Ramos R, et al. A frameshift mutation in NOD2 associated with susceptibility to Crohn's disease. *Nature* 2001; 411:603-6.
31. Hampe J, Cuthbert A, Croucher PJ, Mirza MM, Mascheretti S, Fisher S, et al. Association between insertion mutation in NOD2 gene and Crohn's disease in German and British populations. *Lancet* 2001; 357:1925-8.
32. McFall-Ngai M. Adaptive immunity: care for the community. *Nature* 2007; 445:153.
33. Cucchiara S, Iebba V, Conte MP, Schippa S. The microbiota in inflammatory bowel disease in different age groups. *Dig Dis* 2009; 27:252-8.
34. Ley RE, Hamady M, Lozupone C, Turnbaugh PJ, Ramey RR, Bircher JS, et al. Evolution of mammals and their gut microbes. *Science* 2008; 320:1647-51.
35. Cani PD, Possemiers S, Van de Wiele T, Guiot Y, Everard A, Rottier O, et al. Changes in gut microbiota control inflammation in obese mice through a mechanism involving GLP-2-driven improvement of gut permeability. *Gut* 2009; 58:1091-103.
36. Scalabrino G. Vitamin-regulated cytokines and growth factors in the CNS and elsewhere. *J Neurochem* 2009; 111:1309-26.
37. Grimble RF, Howell WM, O'Reilly G, Turner SJ, Markovic O, Hirrell S, et al. The ability of fish oil to suppress tumor necrosis factor alpha production by peripheral blood mononuclear cells in healthy men is associated with polymorphisms in genes that influence tumor necrosis factor alpha production. *Am J Clin Nutr* 2002; 76:454-9.

38. Lam EK, Tai EK, Koo MW, Wong HP, Wu WK, Yu L, et al. Enhancement of gastric mucosal integrity by *Lactobacillus rhamnosus* GG. *Life Sci* 2007; 80:2128-36.
39. Eisenhardt SU, Thiele JR, Bannasch H, Stark GB, Peter K. C-reactive protein: how conformational changes influence inflammatory properties. *Cell Cycle* 2009; 8:3885-92.
40. Kajander K, Myllyluoma E, Kyronpalo S, Rasmussen M, Sipponen P, Mattila I, et al. Elevated pro-inflammatory and lipotoxic mucosal lipids characterise irritable bowel syndrome. *World J Gastroenterol* 2009; 15:6068-74.
41. Pena JA, Li SY, Wilson PH, Thibodeau SA, Szary AJ, Versalovic J. Genotypic and phenotypic studies of murine intestinal lactobacilli: species differences in mice with and without colitis. *Appl Environ Microbiol* 2004; 70:558-68.
42. Ivanov II, Frutos Rde L, Manel N, Yoshinaga K, Rifkin DB, Sartor RB, et al. Specific microbiota direct the differentiation of IL-17-producing T-helper cells in the mucosa of the small intestine. *Cell Host Microbe* 2008; 4:337-49.
43. Ivanov II, Atarashi K, Manel N, Brodie EL, Shima T, Karaoz U, et al. Induction of intestinal Th17 cells by segmented filamentous bacteria. *Cell* 2009; 139:485-98.
44. Martin FP, Wang Y, Sprenger N, Yap IK, Lundstedt T, Lek P, et al. Probiotic modulation of symbiotic gut microbial-host metabolic interactions in a humanized microbiome mouse model. *Mol Syst Biol* 2008; 4:157.
45. Mackie RI, Sghir A, Gaskins HR. Developmental microbial ecology of the neonatal gastrointestinal tract. *Am J Clin Nutr* 1999; 69:1035-45.
46. Soh SE, Ong DQ, Gerez I, Zhang X, Chollate P, Shek LP, et al. Effect of probiotic supplementation in the first 6 months of life on specific antibody responses to infant Hepatitis B vaccination. *Vaccine*.
47. Isolauri E, Joensuu J, Suomalainen H, Luomala M, Vesikari T. Improved immunogenicity of oral D x RRV reassortant rotavirus vaccine by *Lactobacillus casei* GG. *Vaccine* 1995; 13:310-2.
48. de Vrese M, Rautenberg P, Laue C, Koopmans M, Herremans T, Schrezenmeier J. Probiotic bacteria stimulate virus-specific neutralizing antibodies following a booster polio vaccination. *Eur J Nutr* 2005; 44:406-13.
49. Reiff C, Delday M, Rucklidge G, Reid M, Duncan G, Wohlgenuth S, et al. Balancing inflammatory, lipid and xenobiotic signaling pathways by VSL#3, a biotherapeutic agent, in the treatment of inflammatory bowel disease. *Inflamm Bowel Dis* 2009; 15:1721-36.
50. Reiff C, Kelly D. Inflammatory bowel disease, gut bacteria and probiotic therapy. *Int J Med Microbiol* 300:25-33.
51. Tap J, Mondot S, Levenez F, Pelletier E, Caron C, Furet JP, et al. Towards the human intestinal microbiota phylogenetic core. *Environ Microbiol* 2009; 11:2574-84.
52. Turnbaugh PJ, Hamady M, Yatsunenko T, Cantarel BL, Duncan A, Ley RE, et al. A core gut microbiome in obese and lean twins. *Nature* 2009; 457:480-4.
53. Tschop MH, Hugenholtz P, Karp CL. Getting to the core of the gut microbiome. *Nat Biotechnol* 2009; 27:344-6.
54. Jones SE, Versalovic J. Probiotic *Lactobacillus reuteri* biofilms produce antimicrobial and anti-inflammatory factors. *BMC Microbiol* 2009; 9:35.
55. Peng L, Li ZR, Green RS, Holzman IR, Lin J. Butyrate enhances the intestinal barrier by facilitating tight junction assembly via activation of AMP-activated protein kinase in Caco-2 cell monolayers. *J Nutr* 2009; 139:1619-25.
56. Konstantinov SR, Smidt H, de Vos WM, Bruijns SC, Singh SK, Valence F, et al. S layer protein A of *Lactobacillus acidophilus* NCFM regulates immature dendritic cell and T cell functions. *Proc Natl Acad Sci USA* 2008; 105:19474-9.
57. Sokol H, Pigneur B, Watterlot L, Lakhdari O, Bermudez-Humaran LG, Gratadoux JJ, et al. *Faecalibacterium prausnitzii* is an anti-inflammatory commensal bacterium identified by gut microbiota analysis of Crohn disease patients. *Proc Natl Acad Sci USA* 2008; 105:16731-6.
58. Yasuda E, Serata M, Sako T. Suppressive effect on activation of macrophages by *Lactobacillus casei* strain Shirota genes determining the synthesis of cell wall-associated polysaccharides. *Appl Environ Microbiol* 2008; 74:4746-55.
59. Johnson-Henry KC, Hagen KE, Gordonpour M, Tompkins TA, Sherman PM. Surface-layer protein extracts from *Lactobacillus helveticus* inhibit enterohaemorrhagic *Escherichia coli* O157:H7 adhesion to epithelial cells. *Cell Microbiol* 2007; 9:356-67.
60. Perea Velez M, Verhoeven TL, Draing C, Von Aulock S, Pfitzenmaier M, Geyer A, et al. Functional analysis of D-alanylation of lipoteichoic acid in the probiotic strain *Lactobacillus rhamnosus* GG. *Appl Environ Microbiol* 2007; 73:3595-604.
61. Corr SC, Li Y, Riedel CU, O'Toole PW, Hill C, Gahan CG. Bacteriocin production as a mechanism for the anti-infective activity of *Lactobacillus salivarius* UCC118. *Proc Natl Acad Sci USA* 2007; 104:7617-21.
62. Schlee M, Wehkamp J, Altenhoefer A, Oelschlaeger TA, Stange EF, Fellermann K. Induction of human beta-defensin 2 by the probiotic *Escherichia coli* Nissle 1917 is mediated through flagellin. *Infect Immun* 2007; 75:2399-407.
63. Bergonzelli GE, Granato D, Pridmore RD, Marvin-Guy LF, Donnicola D, Corthesy-Theulaz IE. GroEL of *Lactobacillus johnsonii* La1 (NCC 533) is cell surface associated: potential role in interactions with the host and the gastric pathogen *Helicobacter pylori*. *Infect Immun* 2006; 74:425-34.
64. Makino S, Ikegami S, Kano H, Sashihara T, Sugano H, Horiuchi H, et al. Immunomodulatory effects of polysaccharides produced by *Lactobacillus delbrueckii* ssp. *bulgaricus* OLL1073R-1. *J Dairy Sci* 2006; 89:2873-81.
65. Granato D, Bergonzelli GE, Pridmore RD, Marvin L, Rouvet M, Corthesy-Theulaz IE. Cell surface-associated elongation factor Tu mediates the attachment of *Lactobacillus johnsonii* NCC533 (La1) to human intestinal cells and mucins. *Infect Immun* 2004; 72:2160-9.
66. Kwon HK, Lee CG, So JS, Chae CS, Hwang JS, Sahoo A, et al. Generation of regulatory dendritic cells and CD4⁺Foxp3⁺ T cells by probiotics administration suppresses immune disorders. *Proc Natl Acad Sci USA* 2007; 104:2159-64.
67. Mohan R, Koebnick C, Schildt J, Mueller M, Radke M, Blaut M. Effects of *Bifidobacterium lactis* Bb12 supplementation on body weight, fecal pH, acetate, lactate, calprotectin and IgA in preterm infants. *Pediatr Res* 2008; 64:418-22.
68. Williams HR, Cox IJ, Walker DG, North BV, Patel VM, Marshall SE, et al. Characterization of inflammatory bowel disease with urinary metabolic profiling. *Am J Gastroenterol* 2009; 104:1435-44.
69. Li M, Wang B, Zhang M, Rantalainen M, Wang S, Zhou H, et al. Symbiotic gut microbes modulate human metabolic phenotypes. *Proc Natl Acad Sci USA* 2008; 105:2117-22.
70. Konikoff MR, Denson LA. Role of fecal calprotectin as a biomarker of intestinal inflammation in inflammatory bowel disease. *Inflamm Bowel Dis* 2006; 12:524-34.
71. Rouge C, Piloquet H, Butel MJ, Berger B, Rochat F, Ferraris L, et al. Oral supplementation with probiotics in very-low-birth-weight preterm infants: a randomized, double-blind, placebo-controlled trial. *Am J Clin Nutr* 2009; 89:1828-35.
72. Preidis GA, Versalovic J. Targeting the human microbiome with antibiotics, probiotics and prebiotics: gastroenterology enters the metagenomics era. *Gastroenterology* 2009; 136:2015-31.